by human influence. 1964–1965 and 1986–1987: A comparison. In A. T. Finnamore & S. A. Marshall (eds.), Terrestrial arthropods of peatlands, with particular reference to Canada. Mem.ent. Soc. Can. 169: 61–71.

STRATTON, G. E. & UETZ, G. W. 1986: The inheritance of courtship behavior and its role as a reproductive isolating mechanism in two species of *Schizocosa* wolf spiders (Araneae; Lycosidae). *Evolution* **40**: 129–141.

THALER, K. 1983: Bemerkenswerte Spinnenfunde in Nordtirol (Österreich) und Nachbarländern: Deckennetzspinnen, Linyphiidae (Arachnida: Aranei). Veröff. Mus. Ferdinandeum Innsb. 63: 135–167.

THALER, K. & STEINER, H. M. 1993: Zur epigäischen Spinnenfauna des Stadtgebietes von Wien (Österreich)- nach Aufsammlungen von Prof. Dr W. Kühnelt. Ber. naturw.-med. Ver. Innsbruck 80: 303-310.

Bull. Br. arachnol. Soc. (1995) 10 (2), 74

Nyranytarbus, replacement name for Hemiphrynus Frič, 1901 (Trigonotarbida: Eophrynidae)

Mark S. Harvey

Western Australian Museum, Francis Street, Perth, Western Australia 6000, Australia

and

Paul A. Selden

Department of Geology, University of Manchester, Manchester M13 9PL

The trigonotarbid genus *Hemiphrynus* Frič, 1901 was described for two species of fossil arachnid from the Upper Carboniferous Coal Measures of Nýřany, Czech Republic. These were placed in Araneae, and later Frič (1904) erected a monogeneric family, Hemiphrynidae, for them. Pocock (1910) transferred *Hemiphrynus* to Anthracomartida, and later Petrunkevitch (1953) placed the genus in the family Eophrynidae of the Trigonotarbida.

Since the name *Hemiphrynus* is preoccupied in the Coleoptera by *Hemiphrynus* Horn, 1889, it must be replaced. The name *Hemiphrynus* was also used by Pocock (1902) for an amblypygid genus from South America, which has since been replaced with *Paraphrynus* Moreno, 1940.

During the course of this work, it came to our attention that the date of Frič's (1901) paper is cited differently by various authors; Petrunkevitch (1953), for example, references the same paper twice, once as 1899, and again as 1902! Whilst the cover page of Frič's (1901) work has the data 1899 on it, Frič (1904) refers to it as having been published in 1901. Since this was a work published in parts over a number of years, presumably

the arachnid section appeared after the printing of the title pages. We have therefore taken the date cited by Frič (1904) as the correct year of publication of Frič (1901).

Genus Nyranytarbus, nom. nov.

Hemiphrynus Frič 1901: 57. Junior homonym of Hemiphrynus Horn, 1889 (Coleoptera). Type species Hemiphrynus longipes Frič, 1901, by subsequent designation (Petrunkevitch, 1913: 97).

Etymology: After Nýřany, Czech Republic, the locality from which all known specimens originate; and tarbos, Greek: terror, typical suffix for Trigonotarbida. Gender masculine.

Remarks: Only two species are included in the genus, Nyranytarbus longipes (Frič, 1901), comb. nov. and Nyranytarbus hofmanni (Frič, 1901) comb. nov., both from the Upper Carboniferous Coal Measures of Nýřany, Czech Republic.

References

FRIČ, A. 1901: Fauna der Gaskohle und der Kalksteine der Permformation Böhmens. Vol. IV, part 2. Myriopoda pars II. Arachnoidea: 56–63, pl. 153, 154. Prague.

FRIČ, A. 1904: Palaeozoische Arachniden. 86 pp., 15 pls. Prague. HORN, G. H. 1889: A synopsis of the Halticini of boreal America. Trans. Am. ent. Soc. 16: 163-320.

MORENO, A. 1940: Pedipalpida emendata. Mems Soc.cub.Hist.nat. 'Felipe Poey' 14: 167-168.

PETRUNKEVITCH, A. 1913: A monograph of the terrestrial Palaeozoic Arachnida of North America. Trans. Conn. Acad. Arts Sci. 18: 1-137.

PETRUNKEVITCH, A. 1953: Palaeozoic and Mesozoic Arachnida of Europe. *Mem.geol.Soc.Am.* **53**: xi+128 pp.

POCOCK, R. I. 1902: A contribution to the systematics of the Pedipalpi, Ann. Mag. nat. Hist. (7) 9: 157-165.

POCOCK, R. I. 1910: Notes on the morphology and generic nomenclature of some Carboniferous Arachnida. *Geol.Mag.* (5) 7: 505-512.